Draft version January 10, 2011
Preprint typeset using LaTeX style emulateapj v. 11/10/09

arXiv:1101.1504v1 [astro-ph.SR] 7 Jan 2011

# CAN THE MAXIMUM MASS OF NEUTRON STARS RULE OUT ANY EQUATION OF STATE OF DENSE STELLAR MATTER BEFORE GRAVITY IS WELL UNDERSTOOD?

De-Hua Wen[1,2], Bao-An Li [2,3], Lie-Wen Chen[4]
[1]Department of Physics, South China University of Technology, Guangzhou 510641, P.R. China
[2] Department of Physics and Astronomy, Texas A&M University-Commerce, Commerce, Texas 75429-3011, USA
[3] Department of Applied Physics, Xi'an Jiao Tong University, Xi'an 710049, P.R. China and
[4] Department of Physics, Shanghai Jiao Tong University, Shanghai 200240, P.R. China

*Draft version January 10, 2011*

## ABSTRACT

Probably No! As an example, using soft EOSs consistent with existing terrestrial nuclear laboratory experiments for hybrid neutron stars containing a quark core described with MIT bag model using reasonable parameters, we show that the recently discovered new holder of neutron star maximum mass PSR J1614-2230 of $1.97 \pm 0.04 M_\odot$ can be well described by incorporating a Yukawa gravitational correction that is consistent with existing constraints from neutron-proton and neutron-lead scatterings as well as the spectroscopy of antiproton atoms.

*Subject headings:* hybrid star, hyperon, quark, gravity

## 1. INTRODUCTION

What is gravity? Are there additional spacetime dimensions? These are among the Eleven Science Questions for the New Century identified by the Committee on the Physics of the Universe, US National Research Council (CPUNRC 2003). Interestingly, despite the fact that gravity is the first force discovered in nature, the quest to unify it with other fundamental forces remains elusive because of its apparent weakness at short-distance, see, e.g., refs. (Arkani-Hamed et al. 1998; Pea 2001; Hoyle 2003; Long et al. 2003; Jean et al. 2003; Boehm et al. 2004a,b; Decca et al. 2005). In developing grand unification theories, the conventional inverse-square-law (ISL) of Newtonian gravitational force has to be modified due to either the geometrical effect of the extra spacetime dimensions predicted by string theories and/or the exchange of weakly interacting bosons, such as the neutral spin-1 vector $U$-boson (Fayet 2009), proposed in the super-symmetric extension of the Standard Model, see, e.g., refs. (Adelberger et al. 2003, 2009; Fischbach 1999; Newman 2009; Uzan 2003; Reynaud 2005) for recent reviews. The modified gravity has also been proposed as an explanation for the present period of cosmological acceleration, see, e.g., ref. (DeDeo et al. 2008). The search for evidence of modified gravity is at the forefront of research in several sub-fields of natural sciences including geophysics, nuclear and particle physics, as well as astrophysics and cosmology, see, e.g., refs. (Fujii 1971; Pea 2001; Hoyle 2003; Arkani-Hamed et al. 1998; Long et al. 2003; Adelberger et al. 2009; Kapner et al. 2007; Nesvizhevsky et al. 2008; Kamyshkov et al. 2008; Azam et al. 2008; Geraci et al. 2010; Lucchesi et al. 2010). Various upper limits on the deviation from the ISL has been put forward down to femtometer range. Since the composition of neutron stars are determined mainly by the weak and electromagnetic forces through the $\beta$ equilibrium and charge neutrality conditions while their stability is maintained by the balance of strong and gravitational forces, neutron stars are thus a natural testing ground of grand unification theories of fundamental forces. Moreover, neutron stars are among the densest objects with the strongest gravity in the Universe, making them ideal places to test strong-field predictions of General Relativity (GR) (Psaltis 2008). The masses and radii of neutron stars are solely determined by both the strong-field behavior of gravity and the Equation of State (EOS) of dense stellar matter. However, there is no fundamental reason to choose Einstein's equations over other alternatives and it is known that the GR theory itself may break down at the limit of very strong gravitational fields, see, e.g., ref. (Psaltis 2008) for a comprehensive review. In fact, effects of modified gravity on properties of neutron stars have been under intense investigation. As expected, results of these studies are strongly model dependent, see, e.g., refs. (Germani et al. 2001; Wiseman 2002; Azam et al. 2008; Krivoruchenko et al., 2009; Wen et al. 2009). Nevertheless, it is very interesting to note that alternative gravity theories that have all passed low-field tests but diverge from GR in the strong-field regime predict neutron stars with significantly different properties than their GR counterparts (DeDeo et al. 2003). Moreover, the deviations for neutron star properties from the GR predictions for these theories are larger than the uncertainty due to the poorly known EOS of dense matter in neutron stars. It was also clearly shown that the neutron star maximum mass alone can not distinguish gravity theories (DeDeo et al. 2003). Furthermore, in the endeavor of testing GR theory of gravity using properties of neutron stars, it is known that there is a degeneracy between the matter content and gravity. This degeneracy is tied to the fundamental Strong Equivalence Principle and can only be broken by using at least two independent observables (Yunes et al. 2010).

Recently, using the general relativistic Shapiro delay the mass of PSR J1614-2230 was precisely measured to be $1.97 \pm 0.04 M_\odot$(Demorest et al. 2010), making it the new holder of the maximum mass of neutron stars. Comparing with mass-radius relations predicted from solving the TOV equation using various EOSs within GR theory of gravity, it was shown that the mass of PSR J1614-

2230 can rule out almost all soft EOSs especially those associated with hyperon or boson condensation. While conventional quark stars with soft EOSs are also ruled out by this observation, neutron stars with strongly interacting quark cores are allowed (Demorest et al. 2010; Özel et al. 2010; Lai et al. 2010). It was further shown that a transition to quark matter in neutron star cores can occur at densities comparable to the nuclear saturation density $\rho_0$ only if the quarks are strongly interacting and are color superconducting (Özel et al. 2010). The mass of PSR J1614-2230 was then used to constrain the interacting parameters of quarks. It was also shown that neutron stars with interacting quark clusters in their cores or solid quark stars can be very massive. Using the Lennard-Jones potential for interactions between quark clusters, the mass of the PSR J1614-2230 was used to constrain the number of quarks inside individual quark clusters (Lai et al. 2010). In this work, using soft nuclear EOSs for hybrid stars containing a quark core described by the MIT bag model with reasonable parameters, we show that the mass of PSR J1614-2230 is readily obtained by incorporating a Yukawa gravitational correction that is consistent with existing constraints from terrestrial nuclear laboratory experiments.

## 2. NON-NEWTONIAN GRAVITY AND MODEL EOS OF HYBRID STARS

Fujii (Fujii 1971) first proposed that the non-Newtonian gravity can be described by adding a Yukawa term to the conventional gravitational potential between two objects of mass $m_1$ and $m_2$, i.e.,

$$V(r) = -\frac{Gm_1m_2}{r}(1+\alpha e^{-r/\lambda}), \tag{1}$$

where $\alpha$ is a dimensionless strength parameter, $\lambda$ is the length scale and $G$ is the gravitational constant. In the boson exchange picture, $\alpha = \pm g^2/(4\pi G m_b^2)$ where $\pm$ stands for scalar/vector bosons and $\lambda = 1/\mu$ (in natural units). The $g^2$ and $\mu$ are the boson-baryon coupling constant and the boson mass, respectively. The light and weakly interacting $U$-boson is a favorite candidate mediating the extra interaction (Fayet 2009; Krivoruchenko et al., 2009; Zhu 2007). Similar to the degeneracy between matter content and gravity, there appears to be a duality of incorporating effects of the Yukawa term in either the TOV equation or the input EOS. Nevertheless, according to Fujii (Fujii 1988), the Yukawa term is simply part of the matter system in general relativity. Therefore, only the EOS is modified and the TOV equation remains the same. Within the mean-field approximation, the extra energy density due to the Yukawa term is (Long et al. 2003; Krivoruchenko et al., 2009)

$$\varepsilon_{\rm UB} = \frac{1}{2V}\int \rho(\vec{x}_1)\frac{g^2}{4\pi}\frac{e^{-\mu r}}{r}\rho(\vec{x}_2)d\vec{x}_1 d\vec{x}_2 = \frac{1}{2}\frac{g^2}{\mu^2}\rho^2, \tag{2}$$

where $V$ is the normalization volume, $\rho$ is the baryon number density and $r = |\vec{x}_1 - \vec{x}_2|$. Assuming a constant boson mass independent of the density, one obtains the corresponding addition to the EOS $P_{\rm UB} = \frac{1}{2}\frac{g^2}{\mu^2}\rho^2$, which is just equal to the additional energy density. As it was emphasized by Fujii (Fujii 1988), since the new vector boson contributes to the EOS only through the combination $g^2/\mu^2$, while both the coupling constant g and the mass $\mu$ of the light and weakly interacting bosons are small, the value of $g^2/\mu^2$ can be large. On the other hand, by comparing with the $g^2/\mu^2$ value of the ordinary vector boson $\omega$, Krivoruchenko et al. have pointed out that as long as the $g^2/\mu^2$ value of the U boson is less than about 200 GeV$^{-2}$ the internal structures of both finite nuclei and neutron stars will not change (Krivoruchenko et al., 2009). However, global properties of neutron stars can be significantly modified (Krivoruchenko et al., 2009; Wen et al. 2009) One of the key characteristics of the Yukawa correction is its composition dependence, unlike Einstein's gravity. Thus, ideally one needs to use different coupling constants for various baryons existing in neutron stars. Moreover, to our best knowledge, it is unknown if there is any and what might be the form and strength of the Yukawa term in the hadron-quark mixed phase and the pure quark phase. Nevertheless, instead of introducing more parameters, for the purpose of this exploratory study, we assume that the $P_{\rm UB}$ term is an effective Yukawa correction existing in all phases with the $g$ considered as an averaged coupling constant.



FIG. 1.— (Color online) Model EOSs for hybrid stars with and without the Yukawa contribution using MIT bag constant $B^{1/4}$=170 MeV and $B^{1/4}$=180 MeV, respectively.

Including the Yukawa term the EOS becomes $P = P_0 + P_{\rm UB}$ where $P_0$ is the conventional pressure inside neutron stars. For the latter, we use typical model EOSs for hybrid stars containing a quark core covered by hyperons and leptons. The quark matter is described by the MIT bag model with reasonable parameters widely used in the literature (Chodos et al. 1974; Heinz et al. 1986). The hyperonic EOS is modelled by using an extended isospin- and momentum-dependent effective interaction (MDI (Das et al. 2003)) for the baryon octet with parameters constrained by empirical properties of symmetric nuclear matter, hyper-atoms, and heavy-ion reactions (J. Xu et al. 2010). In particular, the underlying EOS of symmetric nuclear matter is constrained by comparing transport model predictions with data on collective flow and kaon production in relativistic heavy-ion collisions (Danielewicz et al. 2000; Li et al. 2008). Moreover,



FIG. 2.— (Color online) The mass-radius relation of static neutron stars with $B^{1/4}$=170 MeV and various values of $g^2/\mu^2$ in units of GeV$^{-2}$ indicted using numbers above the lines.

the nuclear symmetry energy $E_{sym}(\rho)$ with this interaction is chosen to increase approximately linearly with density (i.e., the MDI interaction with a symmetry energy parameter x=0 (J. Xu et al. 2010)) in agreement with available constraints around and below the saturation density (C. Xu et al. 2010). It is worth noting that this $E_{sym}(\rho)$ is very similar to the well-known APR prediction up to about $5\rho_0$ (Akmal et al. 1998; Xiao et al. 2009). The Gibbs construction was adopted to describe the hadron-quark phase transition (Glendenning 2001). Similar to the previous work in the literature, the hybrid star is divided into the liquid core, inner crust and outer crust from the center to surface. For the inner crust, a parameterized EOS of $P_0 = a + b\epsilon^{4/3}$ is adopted as in refs. (J. Xu et al. 2009). For the outer crust, the BPS EOS (Baym et al. 1971) is adopted. As an example, shown in Fig. 1 are the EOSs for hybrid stars with MIT bag constant $B^{1/4} = 170$ MeV and $B^{1/4} = 180$ MeV, respectively, with and without the Yukawa contribution. The corresponding hadron-quark mixed phase above/below the pure hadron/quark phase covers the density range of $\rho/\rho_0 = 1.31$ to 6.56 and $\rho/\rho_0 = 2.19$ to 8.63, respectively. Including the Yukawa term the EOS stiffens as the value of $g^2/\mu^2$ increases. It is seen that the two sets of parameters, $B^{1/4} = 170$ MeV and $g^2/\mu^2 = 50$ GeV$^{-2}$ or $B^{1/4} = 180$ MeV and $g^2/\mu^2 = 40$ GeV$^{-2}$, lead to approximately the same total pressure. As a reference, the MDI EOS for the $npe\mu$ matter is also shown.

## 3. MAXIMUM MASS OF HYBRID STARS WITH NON-NEWTONIAN GRAVITY

As an example, shown in Fig. 2 is the mass-radius relation of hybrid stars with the bag constant $B^{1/4}$=170 MeV and varying values of $g^2/\mu^2$. First of all, without the Yukawa contribution (black solid line) the maximum stellar mass supported is only about $1.46M_\odot$. Including the Yukawa term, as the EOSs are increasingly stiffened with larger values of $g^2/\mu^2$, the maximum stellar mass increases. With $g^2/\mu^2 = 50$ GeV$^{-2}$ the maximum mass of $1.97M_\odot$ is just in the middle of the measured mass band of PSR J1614-2230. The corresponding radius is about 12.4 km. To see more clearly relative effects of the bag constant B and the Yukawa term, shown in Fig. 3 are the maximum stellar masses as a function of $g^2/\mu^2$ with $B^{1/4} = 170$ MeV and $B^{1/4} = 180$ MeV, respectively. As expected, with $B^{1/4} = 180$ MeV a smaller value of $g^2/\mu^2 = 40$ GeV$^{-2}$ is needed to obtain a maximum mass consistent with the observed mass of PSR J1614-2230.



FIG. 3.— (Color online) The maximum mass of neutron stars as a function of $g^2/\mu^2$ with $B^{1/4}$=170 MeV and $B^{1/4}$=180 MeV, respectively.



FIG. 4.— (Color online) Constraints on the strength and range of the Yukawa term from terrestrial nuclear experiments in comparison with $g^2/\mu^2 \approx 40 - 50GeV^{-2}$.

## 4. COMPARISON WITH TERRESTRIAL CONSTRAINTS ON NON-NEWTONIAN GRAVITY AT SHORT DISTANCE

As mentioned earlier, significant efforts have been devoted to constrain the possible non-Newtonian gravity using terrestrial experiments. These experiments have established a clear trend of increased strength $\alpha$ at shorter length $\lambda$. In the short range down to $\lambda \approx 10^{-14} - 10^{-8}$ m, neutron-proton and neutron-lead scattering data as well as the spectroscopy of antiproton atoms have been used to set upper limits on the value of $g^2/\mu^2$ (or equivalently the $|\alpha|$ $vs$ $\lambda$). It is thus interesting to compare the values of $g^2/\mu^2$ necessary to support the PSR J1614-2230 within the model presented above with the constraints extracted from terrestrial experiments. While the range parameter $\lambda$ is expected to be much larger (smaller) than the radii of finite nuclei (neutron stars), the maximum mass of neutron stars alone is not sufficient to set separate constraints on the values of $\alpha$ and $\lambda$. Shown in Fig. 4 is a comparison with the terrestrial constraints (Kamyshkov et al. 2008; Barbieri et al. 1975; Pokotilovski 2006; Nesvizhevsky et al. 2008) in the $|\alpha|$ *vs* $\lambda$ plane. The straight line is for $g^2/\mu^2 \approx 40-50$ GeV$^{-2}$. It is seen that the values of $g^2/\mu^2$ necessary to describe the maximum mass of PSR J1614-2230 are consistent with the upper limits from the terrestrial experiments.

## 5. CONCLUSIONS

Among all fundamental forces, gravity remains the most uncertain one despite being the first discovered in nature. Neutron stars are natural testing grounds of grand unification theories of fundamental forces. In particular, they are ideal places to test GR predictions at the strong-field limit. Interpretations of observed properties of neutron stars require a comprehensive understanding of both gravity and the EOS of dense stellar matter. Before strong-field gravity is well understood, it is unlikely that the maximum mass of neutron stars alone can rule out any EOS. As an example, using soft nuclear EOSs consistent with existing terrestrial experiments for hybrid stars containing a quark core described by the MIT bag model with reasonable parameters, the maximum mass of PSR J1614-2230 is readily obtained by incorporating the Yukawa gravitational correction that is consistent with existing constraints from terrestrial nuclear laboratory experiments.

We thank W.Z. Jiang, W. G. Newton, A.W. Steiner and Y. Zhang for useful discussions. D.H. Wen is supported in part by the National Natural Science Foundation of China under Grant No.10947023 and the Fundamental Research Funds for the Central University, China under Grant No.2009ZM0193. B.A. Li is supported in part by the US National Science Foundation under grant PHY-0757839, the National Aeronautics and Space Administration under grant NNX11AC41G issued through the Science Mission Directorate and the Texas Coordinating Board of Higher Education under grant No. 003565-0004-2007. L.W. Chen is supported in part by the National Natural Science Foundation of China under Grant Nos. 10675082 and 10975097, MOE of China under project NCET-05-0392, Shanghai Rising-Star Program under Grant No. 06QA14024, the SRF for ROCS, SEM of China, the National Basic Research Program of China (973 Program) under Contract No. 2007CB815004.

## REFERENCES

Adelberger, E. G., Heckel, B. R., & Nelson, A. E. 2003, Annu. Rev. Nucl. Part. Sci., 53, 77

Adelberger, E. G., Gundlach, J. H., Heckel, B. R., Hoedl, S.,& Schlamminger S. 2009, Prog. Part. Nucl. Phys., 62, 102

Akmal, A., Pandharipande, V. R. and Ravenhall, D. G. 1998, Phys. Rev. c, 58, 1804

Arkani-Hamed, N., Dimopoulos, S., & Dvali, G. 1998, Phys Lett. B, 429, 263; Phys. Rev. D, 59, 086004

Azam, M., Sami, M., Unnikrishnan, C. S., & Shiromizu, T. 2008, Phys. Rev. D, 77, 101101

Barbieri, R., & Ericson, T. 1975, Phys. Lett. B, 57, 270

Baym, G., Pethick, C., & Sutherland, P. 1971, ApJ, 170, 299

Boehm, C., Hooper, D., Silk, J., Casse, M. & Paul, J. 2004, Phys. Rev. Lett., 92, 101301

Boehm, C., & Fayet, P. 2004, Nucl. Phys. B, 683, 291

Chodos, A., Jaffe, R. L., Johnson, K., Thorn, C. B., & Weisskopf, V. F. 1974, Phys. Rev. D, 9, 12

Committee on the Physics of the Universe, National Research Council 2003, Connecting Quarks with the Cosmos:Eleven Science Questions for the New Century (The National academies Press)

Danielewicz, P., Lacey, R., & Lynch, W. G. 2000, Science, 298, 1592

Das, C.B., Das Gupta, S., Gale, C., & Li, B. A. 2003, Phys. Rev. D, 67, 034611

Decca, R.S.,López, D., Chan, H. B., Fischbach, E., Krause, D. E., & Jamell, C. R. 2005, Phys. Rev. Lett.94, 240401

DeDeo, S., & Psaltis, D. 2003, Phys. Rev. Lett., 90, 141101

DeDeo, S., & Psaltis, D. 2008, Phys. Rev. D, 78, 064013

Demorest, P., Pennucci, T., Ransom, S., Roberts, M., & Hessels, J. 2010, Nature, 467, 1081

Fayet, P. 2009, Phys. Lett. B, 675, 267

Fischbach, E., & Talmadge, C. L. 1999, The Search for Non-Newtonian Gravity(Springer-Verlag, New York, Inc. )

Fujii, Y. 1971, Nature, 234, 5

Fujii, Y. 1988, in Large Scale Structures of the Universe, page 471-477 (Eds. J. Audouze et al., International Astronomical Union.)

Geraci, A. A., Papp, S. B., & Kitching, J. 2010, Phys. Rev. Lett., 105, 101101

Germani, C., & Maartens, R. 2001, Phys. Rev. D, 64, 124010

Glendenning, N.K. 2001, Phys. Rep., 342, 393

Heinz, U., Subramanian, P. R., Stocker, H., & Greiner, W. 1986, Nucl. Phys., 12, 1237

Hoyle, C. D. 2003, Nature, 421, 899

Jean, P., et al. 2003, A&A, 407, L55

Kamyshkov, Y., Tithof J., & Vysotsky, M. 2008, Phys. Rev. D, 78, 114029

Kapner, D.J., Cook, T. S., Adelberger, E. G., Gundlach, J. H., Heckel, B. R., Hoyle, C. D., & Swanson, H. E. 2007, Phys. Rev. Lett., 98, 021101

Krivoruchenko, M.I., Simkovic, F., & Faessler, A. 2009, Phys. Rev. D, 79, 125023

Lai, X. Y., & Xu, R. X., 2010, arXiv:1011.0526

Li, B. A., Chen, L. W., & Ko, C. M. 2008, Phys. Rep., 464, 113

Long, J. C., et al. 2003, Nature, 421, 922

Lucchesi, D. M., & Person R. 2010, Phys. Rev. Lett., 105, 231103

Nesvizhevsky, V. V., et al., 2008, Phys. Rev. D, 77, 034020

Newman, R. D., Berg, E. C., & Boynton, P. E. 2009, Space Science Review, 148, 175

Özel, F., Psaltis, D., Ransom, S., Demorest, P., & Alford, M, 2010, arXiv:1010.5790v1, ApJL in press.

Pease, R. 2001, Nature, 411, 986

Pokotilovski, Yu. N. 2006, Phys. Atom. Nucl., 68, 924

Psaltis, D. 2008, Living Reviews in Relativity, 11, 9

Reynaud, S., & Jaekel, M. M. 2005, Int. J. Mod. Phys. 20, 2294

Uzan, J. P. 2003, Rev. Mod. Phys., 75, 403

Wen, D. H., Li, B. A., & Chen, L. W., 2009, Phys. Rev. Lett., 103, 211102

Wiseman, T. 2002, Phys. Rev. D, 65, 124007

Xiao, Z.G., et al., 2009, Phys. Rev. Lett. **102**, 062502

Xu, J., Chen, L. W., Ko, C. M., & Li, B. A. 2010, Phys. Rev. C, 81, 055803

Xu, C., Li, B. A., & Chen, L. W. 2010, Phys. Rev. C, 82, 054607

Xu, J., Chen, L. W., Li, B. A., & Ma, H.R. 2009, ApJ, 697, 1549

Yunes, N., Psaltis, D., Ozel F., & Loeb, A. 2010, Phys. Rev. D, 81, 064020

Zhu, S. H. 2007, Phys. Rev. D, 75, 115004